

Department of Pure Mathematics Algebra Comprehensive Examination

May 1994

P. Hoffman and R. Willard

Answer at least six questions, including at least one question from each of the four sections.

I. Group Theory

1. (a) Give an example of a nontrivial group which has no maximal [proper] subgroups.
(b) Consider the following property of a finite group: if H, K are distinct maximal subgroups, then $H \cap K = \{e\}$.
 - i. Describe explicitly all finite *abelian* groups having the stated property.
 - ii. Suppose that G is a finite group having the stated property, H is a maximal subgroup of G , and $H \not\triangleleft G$. Prove that

$$\left| \bigcup_{x \in G} xHx^{-1} \right| \geq \frac{1}{2}|G| + 1.$$

2. (a) Prove that every finite p -group is solvable.
(b) Is there any analogue of (a) concerning groups G for which $|G|$ is divisible by exactly two primes? Explain.
(c) Without using any theorem mentioned in your answer to (b), prove that if p is a prime larger than 3, then every group of order $4p^m$ ($m \geq 1$) is solvable.

II. Ring Theory

3. (a) Give an example (with justification) of a Noetherian integral domain R and an ideal I of R which cannot be generated by fewer than 3 elements.
(b) Prove that, in any Noetherian integral domain, every element is either 0, invertible, or a product of irreducible elements.
(c) Give an example (with justification) of an integral domain R and an element $r \in R$ which is neither 0, invertible, nor a product of irreducible elements of R .
4. (a) Let R be the ring of all matrices $\begin{pmatrix} q & x \\ 0 & y \end{pmatrix}$ where q is a rational number and x, y are real numbers.
 - i. Exhibit a composition series for R_R . *treating R as a left module of itself*
 - ii. Show that ${}_R R$ is not Noetherian.

- (b) Let R be a ring with unit such that
- There exists a simple faithful left R -module;
 - For every $r \in R$ there exists $n \geq 2$ such that $r^n = r$.
- Prove that R is a division ring.

III. Field Theory

- Let F, K be fields with $F \leq K$ and $[K : F]$ finite. Suppose there are only finitely many subfields of K containing F . Prove that $K = F(c)$ for some $c \in K$.
 - Give an example of fields $F \leq K$ such that $[K : F]$ is finite, K is a normal extension of F , but K is not a Galois extension of F .
 - Suppose that F is an algebraically closed field of cardinality 2^{\aleph_0} and characteristic 0. Prove that F is isomorphic to the field \mathbb{C} of complex numbers.
- It is known that the Galois group of $x^4 - x - 1$ over \mathbb{Q} is S_4 . Using this fact, or otherwise, give an example (with justification) of a field K satisfying $\mathbb{Q} \leq K \leq \mathbb{R}$, $[K : \mathbb{Q}] = 4$, and such that there does not exist a field F with $\mathbb{Q} < F < K$.
 - Prove that if K is as in (a), then there is no normal extension E of \mathbb{Q} satisfying $K \leq E$ and $[E : \mathbb{Q}] = 2^n$.
 - If K is as in (a), and if $c \in K \setminus \mathbb{Q}$, does it follow that c cannot be constructed by straightedge and compass? Explain.

IV. Linear Algebra

- Find the smallest $n \in \mathbb{N}$ for which there exist seven matrices A_1, A_2, \dots, A_7 in $\text{Mat}_n(\mathbb{C})$ satisfying both (i) and (ii) below:
 - For each i , the eigenvalues of A_i are exactly the numbers 2, -2 and 3;
 - No two of A_1, \dots, A_7 are similar in $\text{Mat}_n(\mathbb{C})$.
 - Find the smallest $n \in \mathbb{N}$ for which there exist *symmetric* matrices B_1, \dots, B_7 in $\text{Mat}_n(\mathbb{R})$ satisfying both (i) and (ii) below:
 - For each i , the eigenvalues of B_i are exactly the numbers 2, -2 and 3;
 - No two of B_1, \dots, B_7 are orthogonally equivalent.
 - Give an example of two linear operators $\mathbb{C}^n \rightarrow \mathbb{C}^n$ (for some n) which have the same minimal polynomials, same characteristic polynomials, but are not similar in the ring $\text{End}_{\mathbb{C}}(\mathbb{C}^n)$.
 - Let V, W be finite dimensional vector spaces over a field F . Prove that there exists a natural isomorphism $V^* \otimes_F W \rightarrow \text{Hom}_F(V, W)$.
- Let V and W be finitely generated \mathbb{Z} -modules. Let the invariant factors of V be d_1, d_2, \dots, d_k , and let the invariant factors of W be e_1, e_2, \dots, e_l (possibly some are 0). Describe how to compute the invariant factors of $V \otimes_{\mathbb{Z}} W$.